

From: Paul Dulberg Paul_Dulberg@comcast.net 
Subject: Fwd: Walgreens original
Date: November 12, 2022 at 10:47 AM
To: Alphonse Talarico contact@lawofficeofalphonsetalarico.com
Bcc: Paul Dulberg Paul_Dulberg@comcast.net, Tom Kost tkost999@gmail.com

PD

Dear Mr Talarico,

Another recovered document is attached.

We need to supplement the discovery with the recovered files soon because I am adding this as an exhibit to the Draft Baudin complaint and intend to use it in the Mast/Popovich case as well.

FYI - from the time you say send all the recovered documents, it may take me a few days to compile them all together.

The attached document is the original Walgreens RX receipt and when you see it, compared to the 2 different versions currently in the POP bates stamped documents you will notice right away that it was at least a 2 step process to remove the timestamps.

Glad I found this one.

Paul

Begin forwarded message:

From: Paul Dulberg <Paul_Dulberg@comcast.net>
Subject: Walgreens original
Date: October 16, 2022 at 12:20:37 PM CDT
To: Tom Kost <tkost999@gmail.com>

Tom,

Source:
Legal Documents\
Tom\
To Tom\
2019-07-07_For Tom/
2011 Accident/Walgreens\
2011-06-28_Walgreens Prescriptions.pdf

Meta data says original file was Created: April 26, 2019 at 12:24 PM

I believe that date is around when Mike and I got a Mechanical 32 bit hard drive to USB device and transferred old sharp printing files from the dead computers to newer hard drives on Mikes computer.

The transfer was not done for this case but rather was an attempt to save our old artwork and music files.

I had that drive connected to a computer for a few months which includes when you were here in July of 2019 and since it was connected it must have pulled the file when I searched for the word "Walgreens".

I believe this is a unique and earliest version of the Walgreens rx receipt and was probably scanned in back in 2011 before meeting with either Botto or Mast/Popovich.

Let me know what you think.

Paul

**2011-06-28_Walgreens
Prescriptions.pdf**
2.4 MB



YOUR PERSONAL PRESCRIPTION INFORMATION

Your Walgreens Pharmacy Location


3925 W Elm St
McHenry, IL 600504361
(815)363-0722

PATIENT PAUL DULBERG
BIRTH DATE 03/19/70
MEDICATION CEFADROXIL 500MG CAPSULES
QUANTITY 10
DIRECTIONS TAKE 1 CAPSULE BY MOUTH TWICE
DAILY FOR 5 DAYS

DOCTOR DR A. FORD

DRUG DESCRIPTION

PATIENT
ALLERGIES



RED-ORANGE AND WHITE
FRONT: 93 3196

INGREDIENT NAME: CEFADROXIL (sef-a-DROX-ill)

COMMON USES: This medicine is a cephalosporin antibiotic used to treat bacterial infections. This medicine will not treat infections caused by viruses.

BEFORE USING THIS MEDICINE: Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking probenecid. **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions, allergies, pregnancy, or breast-feeding. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. This medicine may be taken on an empty stomach or with food. **STORE THIS MEDICINE** at room temperature, away from heat and light. **TO CLEAR UP YOUR INFECTION COMPLETELY**, continue taking this medicine for the full course of treatment even if you feel better in a few days. Do not miss any doses. **IF YOU MISS A DOSE OF THIS MEDICINE**, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

CAUTIONS: **IF YOU HAVE HAD A SEVERE ALLERGIC REACTION** to a cephalosporin antibiotic (such as Ceclor, Keflex, Ceftin, Duricef) or a penicillin antibiotic (such as amoxicillin, ampicillin), contact your doctor or pharmacist **BEFORE TAKING THIS MEDICINE**. A severe reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine or if a certain medicine is a cephalosporin, contact your doctor or pharmacist. If your symptoms do not improve within a few days or if they become worse, check with your doctor. **IF MODERATE TO SEVERE DIARRHEA OCCURS** during or after treatment with this medicine, check with your doctor or pharmacist. Do not treat it with non-prescription (over-the-counter) medicines. **IF YOU EXPERIENCE** difficulty breathing or tightness of chest, swelling of eyelids, face, or lips, or develop a rash or hives, tell your doctor immediately. Do not take any more of this medicine unless your doctor tells you to do so. Caution should be used in the elderly since they may be more sensitive to the effects of this medicine.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS that

may occur while taking this medicine include dizziness, tiredness, headache, mild diarrhea, or stomach upset. If they continue or are bothersome, check with your doctor. **CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE** if you experience vaginal irritation or discharge, mood or mental changes, or joint pain. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience seizures; yellowing of the eyes or skin; red, blistered, swollen or peeling skin; unusual bruising or bleeding; decreased urination; severe stomach pain or cramps; confusion; or bloody or severe diarrhea. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, severe dizziness, or trouble breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include muscle spasms and seizures.

ADDITIONAL INFORMATION: If your symptoms do not improve within a few days or if they become worse, check with your doctor. **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children.

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

Call your doctor for medical advice about side effects.
You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

YOUR PERSONAL PRESCRIPTION INFORMATION


Your Walgreens Pharmacy Location
3925 W Elm St
McHenry, IL 600504361
(815)363-0722

PATIENT PAUL DULBERG
BIRTH DATE 03/19/70
MEDICATION HYDROCODONE/APAP 10MG/325MG TABS
QUANTITY 20
DIRECTIONS TAKE 1 TABLET BY MOUTH EVERY 6 HOURS AS NEEDED FOR PAIN

DOCTOR DR A. FORD

DRUG DESCRIPTION

PATIENT ALLERGIES



YELLOW

FRONT: WATSON 853

INGREDIENT NAME: HYDROCODONE (hye-droe-KO-done) and ACETAMINOPHEN (a-seat-a-MIN-oh-fen)

COMMON USES: This medicine is a combination of a narcotic and acetaminophen used to relieve moderate to severe pain. Narcotic pain-relievers work by binding to opioid receptors in the brain and spinal cord, and acetaminophen decreases the formation of prostaglandins, therefore reducing pain. This medicine may also be used to treat other conditions as determined by your doctor.

BEFORE USING THIS MEDICINE: Some medicines or medical conditions may interact with this medicine. **INFORM YOUR DOCTOR OR PHARMACIST** of all prescription and over-the-counter medicine that you are taking. **DO NOT TAKE THIS MEDICINE** if you are also taking sodium oxybate. **ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION** may be needed if you are taking other medicines for pain; cimetidine; isoniazid; "blood thinners" (such as warfarin); MAO inhibitors (such as furazolidone, linezolid, moclobemide, phenelzine, procarbazine, selegiline, isocarboxazid, or tranylcypromine); naltrexone; rifampin; or other medicines which cause drowsiness including medicine for sleep (such as zolpidem), tranquilizers, anti-anxiety medicines (such as diazepam), phenothiazines (such as chlorpromazine), tricyclic antidepressants (such as amitriptyline), muscle relaxants, or certain antihistamines (such as diphenhydramine or hydroxyzine). **DO NOT START OR STOP** any medicine without doctor or pharmacist approval. Inform your doctor of any other medical conditions including a history of liver or kidney problems, lung or breathing problems (such as asthma, chronic obstructive pulmonary disease), alcohol use, drug dependency, heart problems (such as slow or irregular heartbeat, very low blood pressure), stomach or bowel problems, gallbladder problems, enlarged prostate, difficulty urinating, seizure disorders, serious head injury or brain disease, spinal problems, low thyroid problems, adrenal gland problems, psychiatric problems, allergies, pregnancy, or breast-feeding. **ROUTINE USE OF THIS MEDICINE IS NOT RECOMMENDED** if you have colitis or diarrhea due to food poisoning. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. Take this medicine by mouth. **THIS MEDICINE MAY BE TAKEN WITH FOOD** if it upsets your stomach, although doing so may decrease its effectiveness. Consult your doctor or pharmacist about alternatives for decreasing nausea (such as antihistamines, or lying down for 1-2 hours with minimal head movement). **STORE THIS MEDICINE** at room temperature in a tightly-closed container, away from heat, and light. **IF YOU MISS A DOSE OF THIS MEDICINE** and you are taking it regularly, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

CAUTIONS: **DO NOT TAKE THIS MEDICINE IF YOU HAVE HAD A SEVERE ALLERGIC REACTION** to codeine, hydrocodone, dihydrocodeine, or oxycodone. A severe allergic reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine or if a certain medicine contains codeine, hydrocodone, dihydrocodeine, or oxycodone, contact your doctor or pharmacist. **DO NOT EXCEED THE RECOMMENDED DOSE** of this medicine. Do not use this medicine more often or for longer than prescribed without checking with your doctor. Exceeding the recommended dose or taking this medicine for longer than prescribed may be habit-forming. If using this medicine for an extended period of time, **DO NOT SUDDENLY STOP** taking this medicine without your doctor's approval. When using for an extended period, this medicine may not work as well and may require different dosing. Talk with your doctor if this medicine stops working well. **KEEP ALL DOCTOR AND LABORATORY APPOINTMENTS** while you are taking this medicine. Laboratory and/or medical tests, including liver, kidney, or lung function tests may be performed to monitor your progress or to check for side effects. This medicine may alter certain lab test results. Make sure that all of your doctors and laboratory personnel know you are taking this medicine. **BEFORE YOU HAVE ANY MEDICAL OR DENTAL TREATMENTS, EMERGENCY CARE, OR SURGERY**, tell the doctor or dentist that you are using this medicine. **THIS MEDICINE MAY CAUSE** drowsiness or dizziness. Using this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. **AVOID ALCOHOLIC BEVERAGES** while taking this medicine. To minimize dizziness or lightheadedness, get up slowly when rising from a seated or lying position. This medicine may cause constipation. To prevent constipation, maintain a diet adequate in fiber, drink plenty of water, and exercise. **THIS MEDICINE CONTAINS ACETAMINOPHEN.** Do not take additional acetaminophen for pain or fever without checking with your doctor or pharmacist.

Ask your pharmacist if you have questions about which medicines contain acetaminophen. If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take this medicine or other pain relievers/fever reducers. Acetaminophen may cause liver damage. Alcohol use combined with this medicine may increase your risk for liver damage. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINE**, either prescription or over-the-counter, check with your doctor or pharmacist. This includes other pain relievers, cough-and-cold medicines, or allergy medicines. **CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY** because they may be more sensitive to the effects of the medicine. **FOR WOMEN: IF YOU PLAN ON BECOMING PREGNANT**, discuss with your doctor the benefits and risks of using this medicine during pregnancy. **AN INGREDIENT IN THIS MEDICINE IS EXCRETED** in breast milk. **IF YOU ARE OR WILL BE BREAST-FEEDING** while you are using this medicine, check with your doctor to discuss the risks to your baby.

POSSIBLE SIDE EFFECTS: **SIDE EFFECTS** that may occur while taking this medicine include nausea, vomiting, constipation, lightheadedness, dizziness, drowsiness, flushing, or vision changes. If they continue or are bothersome, check with your doctor. **CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE** if you experience anxiety, fear, or other mental or mood changes. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience slow or irregular breathing; slow or irregular heartbeat; a change in the amount of urine produced; change or loss in hearing (especially with high doses for long periods); severe drowsiness or dizziness; dark urine; pale stools; or yellowing of the eyes or skin. **AN ALLERGIC REACTION** to this medicine is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, severe dizziness, or trouble breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist. This is not a complete list of all side effects that may occur. If you have questions about side effects, contact your healthcare provider. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include extreme drowsiness, weakness, or fatigue; loss of consciousness; slow, shallow, or abnormal breathing; persistent nausea, vomiting, or stomach pain; slow heartbeat; cold or clammy skin; unusual sweating; and yellowing of the eyes or skin.

ADDITIONAL INFORMATION: If your symptoms do not improve or if they become worse, check with your doctor. **DO NOT SHARE THIS MEDICINE** with others for whom it was not prescribed. **DO NOT USE THIS MEDICINE** for other health conditions. **KEEP THIS MEDICINE** out of the reach of children and pets.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Do not flush unused medications or pour down a sink or drain.

WIC# 957918

KEEP OUT OF REACH OF CHILDREN: STORE IN SAFETY CONTAINER OR SECURE AREA.

